arXiv:1108.4932v1 [hep-ph] 24 Aug 2011

Linear Algebraic Matrix Solution to DGLAP in X Space Leading to Direct, Non-Parametric PDFs extraction From $F_2^{p,d}$ Data

M. Goshtasbpour and M. Zandi
Department of Physics, Shahid Beheshti University, Evin 1983963113, Tehran, Iran

The presented path to PDFs extraction, via global analysis of data, is based on preassumed form of parametric PDFs. In this paper, PDFs are obtained at a chosen Q^2, and at $x_i, i = 1, ..., n$ of the analyzed data points by solving the numerically singular system of decomposition equations of the data points each at a given $(x_i, {Q_{ij}}^2)$ of the data. In each equation, the corresponding data point is decomposed into its pdf components, evolved from the set of unknowns at $(x_i, Q^2), i = 1, ..., n$ to the required $(x_i, {Q_{ij}}^2), i = 1, ..., n, j = 1, ..., n_i$, via our linear algebraic (matrix) solution to DGLAP evolution equations.

INTRODUCTION

Given conditions on parton distribution functions (PDFs), coming from existing measurements in the accelerators, e.g. from a measurement of the structure function F_2 in DIS, through a fixed order (LO, NLO, ...) formula from PQCD for F_2, or a given cross section for pp scattering in a similar manner, can be considered as the initial conditions for the PDFs in the integro-differential DGLAP evolution equations of the same order in perturbation; based on which solving the DGLAP equations generally allows PQCD prediction for processes of interest in accelerators. Thus, the method of solution of DGLAP proposed in the year 2000[1], and developed here can be of serious interest. For simplicity the method is illustrated at LO.

DGLAP evolution equation is of the simple one dimensional form for the non-singlet quark distribution f_{NS}:

$$\frac{\partial f_{NS}(x,t)}{\partial t} = \int_x^1 \frac{dy}{y} P_{qq}(\frac{x}{y}) f_{NS}(y,t) \tag{1}$$

Where

$$t = \frac{33-2n_f}{6} Ln\left(Ln(\frac{Q^2}{\Lambda^2})\right) \tag{2}$$

In the more traditional form of DGLAP the L.H.S. derivative is with respect to $Ln(Q^2)$.

$$\frac{2\pi . dt}{dLn(Q^2)} = \alpha_s(Q^2) = \frac{12\pi}{(33-2n_f)Ln(\frac{Q^2}{\Lambda^2})} \tag{3}$$

Here, the strong coupling constant $\alpha_s(Q^2)$ is rendered in LO in the historical approach[2], using a constant of integration $\Lambda^2 = \Lambda^2_{QCD}$, being in the range $200-400 Mev$, signaling the QCD confinement, where the coupling becomes very strong; e.g. as hinted in LO, (3). We will use the LO formula (3) for the strong coupling constant consistently throughout the calculations later on, avoiding the ambiguities and resulting mistakes that may arise if definition is varied. The present day approach bypasses the ambiguities by directly using experimental determination of $\alpha_s(M_z)$ with no recourse to Λ^2_{QCD}[3, 4].

DGLAP evolution equation has two dimensional form for the singlet quark distribution Σ, coupled to gluons:

$$\frac{\partial}{\partial t}\begin{pmatrix}\Sigma(x,t)\\ g(x,t)\end{pmatrix} = \int_x^1 \frac{dy}{y}\begin{pmatrix}P_{qq}(\frac{x}{y}) & 2n_f P_{qg}(\frac{x}{y})\\ P_{gq}(\frac{x}{y}) & P_{gg}(\frac{x}{y})\end{pmatrix} \cdot \begin{pmatrix}\Sigma(y,t)\\ g(y,t)\end{pmatrix} \tag{4}$$

At LO, where equality of splitting functions (neglecting quark masses), leaves only four different ones, the evolution equation can be written in a simple $(2n+1)$ dimensional form for the n_f quark and ant-iquark flavors coupled through the gluon distribution:

$$\frac{\partial}{\partial t}\begin{pmatrix} q_1(x,t)\\ \vdots \\ q_{2n_f}(x,t)\\ g(x,t)\end{pmatrix} = \int_x^1 \frac{dy}{y}\begin{pmatrix} P_{qq}(\frac{x}{y}) & 0 & \cdots & 0 & P_{qg}(\frac{x}{y})\\ 0 & \ddots & \ddots & \vdots & \vdots\\ \vdots & \ddots & \ddots & 0 & \vdots\\ 0 & \cdots & 0 & P_{qq}(\frac{x}{y}) & P_{qg}(\frac{x}{y})\\ P_{gq}(\frac{x}{y}) & \cdots & \cdots & P_{gq}(\frac{x}{y}) & P_{gg}(\frac{x}{y})\end{pmatrix} \cdot \begin{pmatrix} q_1(y,t)\\ \vdots\\ q_{2n_f}(y,t)\\ g(y,t)\end{pmatrix} \tag{5}$$

N-DIMENSIONAL DISCRETIZATION OF X-SPACE OF THE DATA POINTS AND PDFS.

Analytical solutions to DGLAP exploits nearness of the discrete data points of a given set, e.g. of the structure functions, to the Bjorken x_i,

$$x_i = (x_1)^i, \qquad 0 < ... < x_{i+1} < x_i < ... < x_1 < 1, i = 1, ..., n. \tag{6}$$

Numerical solutions can simply use the exact Bjorken $x_i, i = 1, ..., n$, of the data set. In either case, a discrete set of n basis vectors for the PDFs is defined on each x_i. Thus, the splitting functions operating on this basis (space) can be calculated as $n \times n$ matrices.

The discrete basis for the PDFs are used in the convolution integrals on the right side of the DGLAP equation to produce the matrix form of the kernel K, which in DIS data analysis, is either the splitting functions for DGLAP, or the coefficient functions of the hadronic structure function. The convolution integrals are of the generic form:

$$P \otimes f = \int_x^1 \frac{dy}{y} K(\frac{x}{y}) f(y, Q^2) = \int_x^1 \frac{dy}{y} K(y) f(\frac{x}{y}, Q^2), \tag{7}$$

in which, for the simplest parton distribution function, e.g. non-singlet (NS) $f(x, Q^2) = f_{NS}(x, Q^2)$, the discrete basis is an $n-$tuple, whose ith element is $f(x_i, Q^2)$. As the convolution integral is made a discrete sum, $P_{ij}, i, j = q, g$ or $C_k, k = q, g$ within the kernel K become lower triangular (l.t.), due to the limits of integration of the resulting sum; and banded (bdd), due to divisions of x_i in (1). The resulting algebraic simplicity of the l.t. bdd matrices is an essential element for having analytical solution of DGLAP as the basis of the analysis. Evidently, numerical solution working with exact $x_i, i = 1, ..., n$ of the data neither works with, nor requires bandedness.

Illustration of the method by an example

To calculate the matrix form of a splitting function, e.g. P_{qq} in (1), a combination of discrete finite difference approximation of the unknown PDFs and continuous integration of the known splitting functions is mixed in our method of evaluation of the convolution, as follows. Then, $(P_{qq})_{ik}$ is extracted from the coefficients of $f_k = f(x_k, t)$. At LO, P_{qq} is given in the following equivalent forms[5]:

$$P_{qq} = \frac{4}{3}\left(\frac{1+y^2}{1-y}\right)_+ = \frac{4}{3}\left[\frac{1+y^2}{(1-y)_+} + \frac{3}{2}\delta(1-y)\right]. \tag{8}$$

Given the defnition of "+" regularization, aimed at removal of the infinity at $y = 1$ in the kernel P_{qq} in the context of the convolution integral[6]:

$$\int_z^1 dx f(x)\left[\frac{g(x)}{1-x}\right]_+ = \int_z^1 dx \frac{f(x) - f(1)}{1-x} g(x) - f(1)\int_0^z dx \frac{g(x)}{1-x}, \tag{9}$$

The convolution (7) becomes:

$$I = \int_x^1 \frac{dy}{y} y . \frac{4}{3}\left(\frac{1+y^2}{1-y}\right)_+ f(\frac{x}{y}) = \int_x^1 dy \frac{4}{3}\left(\frac{1+y^2}{1-y}\right) f(\frac{x}{y}) - f(x)\int_0^1 dy \frac{4}{3}\left(\frac{1+y^2}{1-y}\right) = I_1 + I_2, \tag{10}$$

where the extra y in the left integrand is put in to work with the more useful form, the momentum distribution. Note that the DGLAP equation can be written for momentum distribution, i.e, in $f(x) \to xf(x)$, by putting an extra x on both sides, and a factor of $y.\frac{1}{y}$ within the integral. Different techniques may be used for doing the integration in DGLAP. Here, we use only integration by parts:

$$u = f(\frac{x}{y}) \Rightarrow du = \frac{df(\frac{x}{y})}{dy} dy, \; dv = \frac{4}{3}(\frac{1+y^2}{1-y}) dy \Rightarrow v = \int dv;$$

So,

$$I_1 = f(x)v(1) - f(1)v(x) - \int_x^1 v(y)\frac{df(\frac{x}{y})}{dy} dy = f(x)v(1) + \int_x^1 v(\frac{x}{y})\frac{df(x)}{dy} dy,$$
$$I_2 = -f(x)[v(1) - v(0)]. \tag{11}$$

$$\Rightarrow I = f(x)v(0) + \int_x^1 v(\frac{x}{y})\frac{df(x)}{dy} dy.$$

Here and later, when convenient, the variable change $\frac{x}{y} \to y$ is used. Bringing in a finite difference approximation of the differentials:

$$\begin{aligned} I &= v(0)f(x_i) + \sum_{k=1}^{i} \int_{x_k}^{x_{k-1}} v(x_i/y)\frac{f_k - f_{k-1}}{x_k - x_{k-1}} dy \\ &= v(0)f(x_i) + \int_{x_i}^{x_{i-1}} v(x_i/y)\frac{f_i - f_{i-1}}{x_i - x_{i-1}} dy + \\ &\sum_{k=1}^{i-1} \int_{x_k}^{x_{k-1}} v(x_i/y)\frac{f_k - f_{k-1}}{x_k - x_{k-1}} dy \qquad (12) \\ &= v(0)f(x_i) + \frac{f_i}{x_i - x_{i-1}} \int_{x_i}^{x_{i-1}} v(x_i/y) dy - \\ &\frac{f_{i-1}}{x_i - x_{i-1}} \int_{x_i}^{x_{i-1}} v(x_i/y) dy + \sum_{k=1}^{i-1} \int_{x_k}^{x_{k-1}} v(x_i/y)\frac{f_k - f_{k-1}}{x_k - x_{k-1}} dy. \end{aligned}$$

Now, a bdd l.t. P_{qq} can be read (extracted) from the coefficients of f_k .

$$P_{qq} = \begin{cases} (P_{qq})_{ii} = v(0) + \dfrac{1}{x_i - x_{i-1}} \displaystyle\int_{x_i}^{x_{i-1}} v(x_i/y)dy, \\ (P_{qq})_{ik} = \dfrac{1}{x_k - x_{k-1}} \displaystyle\int_{x_k}^{x_{k-1}} v(x_i/y)dy- \\ \dfrac{1}{x_{k+1} - x_k} \displaystyle\int_{x_{k+1}}^{x_k} v(x_i/y)dy. \end{cases} \quad (13)$$

The other three kernels are essentially obtained in a similar manner. In addition, construction of the matrix coefficient functions in $\overline{MS}$ which are taken from[2], follows closely the path of construction of splitting functions, resulting in commuting, bdd, l.t. or non-commuting, non-bdd, l.t. matrices, respectively, in analytical or more general numerical approaches.

LINEAR ALGEBRAIC (MATRIX) FOUNDATIONS OF THE NUMERICAL SOLUTION OF DGLAP IN X-SPACE

In this section, the numerical solution of the DGLAP equations (1), (4) and (5) is explained in detail. The DGLAP equations are of the generic form:

$$\frac{df_K(x,t)}{dt} = \int_x^1 \frac{dy}{y} K(\frac{x}{y}) f_K(y,t), \quad (14)$$

Where $K = P_{NS}, P_S, P_F$ is one dimensional in non-singlet, two dimensional in singlet and $(2n_f + 1)$ dimensional in flavor kernels, respectively.

Having the matrix forms of the above kernels from section (2), the matrix form of DGLAP equations reads as:

$$\frac{df_i^K(t)}{dt} = \sum_{j=1}^{i} K_{ij} f_j^K(t) \Leftrightarrow \mathbf{f}^K(t) = \mathbf{K}.\mathbf{f}^K(t); \quad (15)$$

Since the kernel K being independent of the variable t, leads the following solution to the matrix equation (15) for finite evolution from t_0 to t:

$$\mathbf{f}^K(t) = e^{\mathbf{K}(t-t_0)}.\mathbf{f}^K(t_0) \equiv \mathbf{E}_t^K.\mathbf{f}^K(t_0). \quad (16)$$

Therefore, the finite evolution machinery for the DGLAP, is constructed with the final $(t - t_0)$ dependent $\mathbf{E}_t^K$ matrix.

To determine the matrix form of $\mathbf{E}_t^K = e^{\mathbf{K}(t_0-t)}$ in analytical framework of[1], as mentioned in the previous sections, the choice of $x_i = x_1^i, i = 1, \cdots, n$ is mandatory to have commuting bdd splitting functions matrices, but the experimental data are not always match these x points, so interpolation techniques should be used to shift the data points to the desired places.

On the other hand, given the numeric form of the LO kernels P_{NS}, P_S and P_F to a software such as *Mathematica* (using the MatrixExp command), immediately gives the matrix form of the exponential in the solution (16). Using numerical algorithms to approximate the infinite series of the exponential has the advantage of being needless of the commuting bdd matrices, so, ones could chose the experimental x to discrete the x space.

COMPARISON OF THE EVOLUTION MACHINE RESULTS WITH MSTW AND CTEQ

The machinery thus far has brought us to the point that we may take a discrete set of input PDFs at x-points patterned according to (6) at t_0 from some present standard LO set, such as MSTW2008, evolve it ourselves to t, and then compare our results with their PDFs at t. This is actually done via an $n = 100$ (6) input taken at $Q_0^2 = 10Gev^2$ from MSTW (green curves). Evolution needs the value of α_s which, in the line of the discussion of equation (3), was handled using Λ_{QCD} consistently. The excellent quality of match of our solutions outputs at $Q^2 = 100Gev^2$ (red curves) and MSTW PDFs (blue curves) can be seen in Fig.(1-I) for non-singlet - singlet, and in Fig.(1-II) for the flavors.

Definitions of 5 flavor non-singlets and the singlet are:

$$\begin{aligned} &q_3^{NS} = (u + \bar{u}) - (d + \bar{d}), \\ &q_8^{NS} = (u + \bar{u}) + (d + \bar{d}) - 2(s + \bar{s}), \\ &q_{15}^{NS} = (u + \bar{u}) + (d + \bar{d}) + (s + \bar{s}) - 3(c + \bar{c}), \\ &q_{24}^{NS} = (u + \bar{u}) + (d + \bar{d}) + (s + \bar{s}) + (c + \bar{c}) - 4(b + \bar{b}), \\ &\Sigma = (u + \bar{u}) + (d + \bar{d}) + (s + \bar{s}) + (c + \bar{c}) + (b + \bar{b}). \end{aligned} \quad (17)$$

A surprise is that, we assume complete $SU_f(5)$(flavor) symmetry with our up to q_{24}^{NS}, or flavor DGLAP Kernel (5). Then, an $n = 100$ point fit has such a good match with MSTW2008 that says they should be having such an $SU_f(5)$ symmetry assumption as well. The only source of asymmetry in both our procedures seems to be the non- symmetric PDFs inputs to the totally symmetric evolution process, resulting in the asymmetric outputs. A similar comparison is shown in figure(1-III) with the older LO CTEQ5L set. It is interesting to note that the excellent match, points to their $SU_f(5)$ symmetry assumption as well.

There is only a final step needed to have the detailed match observed in the figures. While solving DGLAP, flavor number changes should be accommodated every time a changing Q^2 crosses a massive quarks m_q^2. Accordingly, a simple method of continuous evolution across stages with breaks at $Q^2 = m_q^2$ was conceived, which may be called zero-mass scheme. A more involved zero-mass scheme is found in MSTW [4]. But there is no apparent difference for our LO comparison! In both, as mentioned there, the name zero-mass is misleading, as mass appears in the boundary conditions. Continuity of PDFs is kept via the boundary conditions of the DGLAP at breaks at $Q^2 = m_q^2$. Specifically, a break takes place if evolution of the PDFs, during increase of $t(Q^2)$ (2), from n_f flavors at t_0, crosses a $t_q \equiv t(m_q^2)$, of a massive quark h, to reach t with $n_f + 1$ flavors. Then, solving DGLAP for finite evolutions $[t_0, t]$ is divided into two stages of $[t_0, t_q]$ and$[t_q, t]$. Furthermore, there are two sets of inputs, the first set has n_f flavors at t_0, the second set has $(n_f + 1)$ flavors at t_q. In addition, the second set of inputs is the first set of outputs with an extra input PDF for the heavy quark, $h(m_q^2) \equiv 0$. Specifically, in our figures, h= b-quark with MSTW $m_b = 4.75 Gev$, and CTEQ $m_b = 4.5 Gev$.

EXTRACTION OF PDFS FROM ${F_2}^{p,d}$ DATA

Beginning with LO structure function F_2 symbolically as:

$$F_2 = x(1 + \frac{\alpha_s}{2\pi} C_q) \otimes \bar{F}_2 + x\frac{\alpha_s}{2pi} C_g \otimes (\sum_f e_f^2) g. \quad (18)$$

with the quark structure function in term of $n_f = 5$ flavors NS-S, with usual definitions (17)[7] ,

$$\bar{F}_2^p = \frac{1}{90}(22\Sigma + 3q_{(24)} - 5q_{(15)} + 5q_{(8)} + 15q_{(3)}), \quad (19)$$

$$\bar{F}_2^d = \frac{F_2^p + F_2^n}{2} = \frac{1}{90}(22\Sigma + 3q_{(24)} - 5q_{(15)} + 5q_{(8)}).$$

Now, we are ready to actualize the program explained at the onset. Each $F_2^{p,d}$ data point at a given (x_i, Q_{ij}^2) of an experiment, e.g. BCDMS[8], is decomposed into its partonic components via (18) and (19), evolved by our solutions of DGLAP, from a set of unknown parton distributions at (x_i, Q^2) to be solved for by the linear system of equations for all the data and the variable flavor number (VFN) constraints. Our data equations come from minimization of χ^2 with respect to the unknowns, with the usually defined[4]:

$$\chi^2 = \sum_{i=1}^{N_{pts.}} \frac{(F_{2i}^L - F_{2i}^R)^2}{\sigma_i^2} \quad (20)$$

where F_{2i}^L is the value of data point on the left side of (18) with its quadratically calculated total error being σ_i, and F_{2i}^R is its LO decomposition corresponding to the right side of (18). For the range of Q^2 used, the VFN constraint equations are set at the masses of c and b quarks, considered as boundary conditions (b.c.) between regions of evolution in Q^2 where n_f respectively crosses 3- 4, and 4-5. Constraints equate the singlet, Σ, and the $(n_f^2 - 1)$ -multiplet NS of the $SU(n_f)$ flavor symmetry above each b.c.; thus, dropping the flavor number, n_f, by 1 at the b.c. The constraint equations, together with consistent evolution, keep the continuity of the PDFs while the flavor numbers changes at the b.c.

RESULTS

As mentioned in the previous section, each data point serves as a constraint equation on PDFs. A measurement of the structure function F_2 in DIS, through a fixed order (LO, NLO,...) formula from PQCD for F_2, or a given cross section for pp scattering in a similar manner, can be considered as such a constraint equation. The integro-differential DGLAP evolution equations of the same order in perturbation may then be solved and used to tie all the data points in an over determined and numerically singular system of linear equations to be solved via singular value decomposition (SVD)[9] together with other constraint equations on the PDFs coming from variable flavor numbers (VFN) pertinent to the Q^2 range of the data. Here, the presented method of solution of DGLAP, becomes of serious interest. The presentation here is done at LO, for the simplest example of DIS data, $F_2^{p,d}$ from BCDMS collaboration[8] ; however, it is treated in general terms for other applications and applicable to be extended through PDF Global Data Analysis as well as higher orders of perturbation.

Solutions for the PDFs are presented in figure(2). The error analysis comes from error propagation equation[10], applied on the output of the solved system of equations. For simplicity, we assumed no correlation among experimental errors. Total number of used data points, for proton and deuteron respectively, are $n_p = 112$ and $n_d = 96$. Number of data equations (same as the number of unknowns) obtained from χ^2 are 6(PDFs) ×11(x-points), and those obtained from b.c. are 2(b.c.)×11(x-points) and $\chi^2 / N_{pts.} = 0.93$. Noting that MSTW2008LO PDFs are coming from global fits with the total number of 2615 data points from 36 data sets.

CONCLUSION

We have developed in detail, analytical linear algebraic solution to DGLAP, with its numerical presentation, employed it in doing non-parametric data analysis; i.e. obtaining PDFs directly from data points without a pre-assumed parametric form. Our results for finite LO evolution match those of the simply available standard LO sets: MSTW2008 and CTEQ5L. In addition, the obtained parton distributions agree with the sophisticated



FIG. 1. (a-I, II, III) LO Parton distributions extracted from MSTW2008 Ns-S, Flv. And CTEQ5L Ns-S respectively at $Q^2 = 10Gev^2$. (b-I, II, III) Parton distributions at $Q^2 = 100Gev^2$ (blue curves) in comparison with its corresponding evolved distributions from $Q_0^2 = 10Gev^2$ to $Q^2 = 100Gev^2$ using the mentioned technique (red dashed curves).



FIG. 2. Reasonable matches among extracted PDFs (red dots) from BCDMS $F_2^{p,d}$ DIS data in the range of $x = 0.07$ to $x = 0.75$ in comparison with MSTW2008LO global fit distributions (blue curves) at fixed $Q^2 = 37.5 Gev^2$.

MSTW and CTEQ global data analysis.

ACKNOWLEDGMENTS

We would like to mention that SBU physics students, notably S. A. Shafiei Darabi with his M.Sc. thesis and A. Mollabashi with his B.S. project, took part in building an edifice which this paper is a part of.

[1] Goshtasbpour, M. and Ratcliffe, P. G., 14th International Spin Physics Symposium, Osaka, Japan, Oct. 16-21, 2000, P. 879; Ratcliffe, P. G., Phys. Rev. D63, 116004 (2001), arXiv: hep-ph/ 0012376; S.A. Shafiei, M.Sc. Thesis, A Study of DIS and a Novel Solution to DGLAP Evolution Equation, Shahid Beheshti University, 2000-2002.

[2] R.K. Ellis, W.J. Stirling, B.R. Webber, *QCD and Collider Physics*, Cambridge university press (1996).

[3] Ibid, p.38.

[4] A.D. Martin, W.J. Stirling, R.S. Thorne, G. Watt, *Eur.Phys.J.* **C**63:189-285,2009.

[5] Guid Alterelli, Parons in QCD, Table 2, p.40.

[6] G.Sterman, et all, *Handbook of perturbative QCD* ,CTEQ Collaboration(1999), chapter3.

[7] G. Soyez, *Phys.Rev.* D**71**,076001 (2005).

[8] Benvenuti et al., *phys. Lett.* B**223**,485 (1989); *phys. Lett.* B**237**,529 (1990).

[9] W.H. Press, S.A. Teukolsky, W.T. Vetterling, B.P. Flannery *Numerical Recipes in C: The Art of Scientific Computing* , Cambridge university press (1992).

[10] Bevington P.R., Robinson D.K, *Data reduction and error analysis for physical sciences*,McGraw-Hill(2003).